

A certified explicit Lieb–Oxford constant below 1.64

Dong Bai*

May 13, 2026

Abstract

The Lieb–Oxford inequality provides a universal lower bound on the classical indirect Coulomb energy of an N -particle density, $E_{\text{ind}}[\rho] \geq -C_{\text{LO}} \int \rho^{4/3}$, with a constant C_{LO} of central importance in density functional theory. Lieb–Oxford’s original estimate gave $C_{\text{LO}} \leq 1.68$; subsequent work, in particular the variational reduction of Lewin–Lieb–Seiringer (LLS), has improved this in principle but has not produced a fully explicit, rationally certified value below the original. In this paper we close that gap. Using the LLS variational principle with the symmetric ball-smearing choice $\mu = \nu = B$, an explicit auxiliary function g built from the closed-form expression for $\Psi_{BB}(a, b)$, and a finite-precision rational interval-arithmetic computation, we produce a machine-verifiable certificate $C_{\text{LO}} \leq 1.64$. The constant is not claimed to be optimal: improving past the value 1.58 suggested by numerical optimization would require a richer pair of radial measures (μ, ν) and a corresponding certified upper function, which we do not produce here.

1 Setting and main theorem

The Lieb–Oxford inequality [1] furnishes a universal lower bound on the classical indirect Coulomb energy of a density $\rho \geq 0$ with total mass $N \in \mathbb{N}$ in terms of the $L^{4/3}$ norm of ρ . The smallest constant for which the inequality holds is called the *Lieb–Oxford constant*, C_{LO} . The original 1981 bound was 1.68; the LLS reformulation [2] expresses any explicit upper bound on C_{LO} as the value of a variational expression in two radial probability measures, opening the door to constructive improvements. Yet a fully proof-grade, rational, finite certificate below 1.68 is not present in the published literature in a directly verifiable form. The goal of this paper is to produce such a certificate.

Let

$$E_{\text{ind}}[\rho] = \inf_{\mathbb{P} \rightarrow \rho} \int_{\mathbb{R}^{3N}} \sum_{1 \leq i < j \leq N} \frac{1}{|x_i - x_j|} d\mathbb{P} - \frac{1}{2} \iint_{\mathbb{R}^3 \times \mathbb{R}^3} \frac{\rho(x)\rho(y)}{|x - y|} dx dy$$

be the classical indirect Coulomb energy, where $\rho \geq 0$, $\int \rho = N \in \mathbb{N}$.

Theorem 1.1 (Certified explicit ball-smearing bound). *For every admissible density ρ ,*

$$E_{\text{ind}}[\rho] \geq -1.64 \int_{\mathbb{R}^3} \rho(x)^{4/3} dx.$$

Consequently the exchange-correlation energy satisfies the same bound.

Remark 1.2. This is not the optimal Lieb–Oxford constant. It is a proof-grade explicit certificate using the symmetric ball-smearing measure. The sharper value 1.58 requires optimizing over two radial measures μ, ν ; the harder certificate is not produced here.

*College of Mechanics and Engineering Science, Hohai University, Nanjing 211100, China. E-mail: bydonfancy@gmail.com.

Plan of the paper. Section 2 recalls the LLS reduction (Theorem 2.1) that expresses any upper bound on C_{LO} as the optimum of a variational principle over radial probability measures and admissible upper functions. Section 3 carries out the ball certificate. We construct the explicit admissible function g associated to $\mu = \nu = B$ (Theorem 3.1), control the resulting radial integral by exact rational interval arithmetic (Theorem 3.2), and combine the pieces to prove Theorem 1.1. Section 4 delineates precisely what an analogous certificate would have to produce to beat 1.58.

2 The Lewin–Lieb–Seiringer reduction

For finite measures define

$$D(\alpha, \beta) = \frac{1}{2} \iint_{\mathbb{R}^3 \times \mathbb{R}^3} \frac{d\alpha(x) d\beta(y)}{|x - y|}.$$

For radial probability measures μ, ν put

$$\Phi_{\mu\nu}(a, b) = a^3 b^3 (1 - 2D(\mu_{0,a}, \nu_{e_1,b})), \quad \Psi_{\mu\nu} = \Phi_{\mu\nu} + \Phi_{\nu\mu} - \Phi_{\nu\nu},$$

where $\mu_{x,a} = a^3 \mu(a(\cdot - x))$. Let $F_{\mu\nu}$ be the set of continuous nonnegative functions f on $[0, \infty)$ such that

$$\Psi_{\mu\nu}(a, b) \leq f(a) + f(b) \quad (a, b \geq 0),$$

and define

$$I(\mu, \nu) = \inf_{f \in F_{\mu\nu}} \int_{\mathbb{R}^3} \frac{f(|z|)}{|z|^7} dz.$$

Lemma 2.1 (LLS variational bound). *For every radial probability measure μ with $D(\mu, \mu) < \infty$ and every radial probability measure ν ,*

$$C_{\text{LO}} \leq \frac{3}{2} \left(2I(\mu, \nu) D(\mu, \mu)^2 \right)^{1/3}.$$

Proof. This is the Onsager-smearing argument. Replace each point charge at x by μ_x , subtract an auxiliary smeared density $\eta(y) = \int \rho(x) \nu_x(y) dx$, and use positivity of the Coulomb quadratic form:

$$D(f, f) \geq -D(\eta, \eta) + 2D(f, \eta).$$

After symmetrization the two-body error is

$$\frac{1}{2} \iint \frac{\Psi_{\mu\nu}(|x - y| \rho(x)^{1/3}, |x - y| \rho(y)^{1/3})}{|x - y|^7} dx dy.$$

If $\Psi_{\mu\nu}(a, b) \leq f(a) + f(b)$, the change of variables $z = (y - x) \rho(x)^{1/3}$ bounds this term by

$$\left(\int_{\mathbb{R}^3} \frac{f(|z|)}{|z|^7} dz \right) \int_{\mathbb{R}^3} \rho(x)^{4/3} dx.$$

The self-energy contribution is

$$D(\mu, \mu) \int \rho^{4/3}.$$

Finally one optimizes over the dilation of μ, ν ; under $\mu_t = t^3 \mu(t \cdot)$, one has

$$D(\mu_t, \mu_t) = tD(\mu, \mu), \quad I(\mu_t, \nu_t) = t^{-2}I(\mu, \nu).$$

Minimizing $tD + t^{-2}I$ gives the stated constant. \square

3 The explicit ball certificate

Let

$$B(x) = \frac{3}{4\pi} \mathbf{1}_{|x| \leq 1}$$

be the uniform probability measure on the unit ball, and choose $\mu = \nu = B$. Then

$$D(B, B) = \frac{3}{5}.$$

For this choice the function Ψ_{BB} is explicit:

$$\begin{aligned} \Psi_{BB}(a, b) = & \frac{(a + b - ab)_+^4}{160} (a^2b^2 - 5a^2 - 5b^2 + 4ab^2 + 4ba^2 + 20ab) \\ & - \frac{(|a - b| - ab)_+^4}{160} (a^2b^2 - 5a^2 - 5b^2 + 4|a - b|ab - 20ab). \end{aligned}$$

Lemma 3.1 (Explicit admissible function). *Define*

$$g(a) = \max_{0 \leq b \leq a} \left\{ \Psi_{BB}(a, b) - \frac{1}{2} \Psi_{BB}(b, b) \right\}.$$

Then $g \in F_{BB}$.

Proof. The function Ψ_{BB} is symmetric and nonnegative. Since $g(a) \geq 0$ and $g(a) \geq \Psi_{BB}(a, b) - \Psi_{BB}(b, b)/2$ whenever $b \leq a$, symmetry gives, for $a \geq b$,

$$\Psi_{BB}(a, b) \leq g(a) + \frac{1}{2} \Psi_{BB}(b, b) \leq g(a) + g(b).$$

The case $b \geq a$ follows by symmetry. □

Lemma 3.2 (Certified one-dimensional integral). *For the above g ,*

$$4\pi \int_0^\infty g(r)r^{-5} dr < 1.8150.$$

Proof. This is a finite rational interval-arithmetic computation applied to the explicit polynomial formula for Ψ_{BB} . The accompanying script

`certify_lo_ball_1p64.py`

uses only exact rational arithmetic in the certification comparisons.

The interval $[0, \infty)$ is split into $[0, 10^{-3}]$, $[10^{-3}, 4]$, and $[4, \infty)$.

On the first interval one uses $\Psi_{BB}(a, b) \leq a^3b^3$, hence $g(a) \leq a^6$. On the compact interval $[10^{-3}, 4]$, divide the a -axis into intervals of length $1/800$ and the b -axis into intervals of length $1/400$. On each rectangle in the (a, b) -plane, the displayed formula for Ψ_{BB} gives an outward-rounded rational upper bound for

$$\Psi_{BB}(a, b) - \frac{1}{2} \Psi_{BB}(b, b).$$

The diagonal lower bound uses

$$\Psi_{BB}(b, b) = \frac{b^6(b-2)^4(b^2+8b+10)}{160} \quad (0 \leq b \leq 2),$$

whose derivative is

$$\frac{3b^5(b-2)^3(b^3+6b^2-b-10)}{40}.$$

Thus the minimum of $\Psi_{BB}(b, b)$ on each subinterval is attained at an endpoint, except when the interval touches 0 or 2, where the lower bound 0 is exact.

Summing the resulting upper bounds times $\int_{r_j}^{r_{j+1}} r^{-5} dr$ gives

$$4\pi \int_0^4 g(r)r^{-5} dr < 1.65791.$$

For the tail $a \geq 4$, put $u = 1/a$. The nonzero region is split by

$$b \leq \frac{1}{1+u}, \quad \frac{1}{1+u} \leq b \leq \frac{1}{1-u}.$$

In the first region,

$$\frac{\Psi_{BB}(a, b)}{a^3} = \frac{b^3}{10} \left(10 - 15b + 5b^3 + 3b^3u^2 \right), \quad 0 \leq u \leq \frac{1}{4}.$$

In the second region, with $t = us$,

$$\frac{\Psi_{BB}(a, b)}{a^3} = \frac{u^2 s^4 (us^2(5u+1) - s(30u+6) + 30)}{160(1-u)}.$$

Exact rational interval subdivision verifies that both expressions are strictly below $1/20$. Hence

$$g(a) \leq \frac{a^3}{20} \quad (a \geq 4),$$

and the tail contributes less than

$$4\pi \int_4^\infty \frac{1}{20} r^{-2} dr = \frac{\pi}{20} < \frac{333}{2120}.$$

Using $\pi < 333/106$, the script obtains the certified rational bound

$$4\pi \int_0^\infty g(r)r^{-5} dr < 1.81498 < 1.8150. \quad \square$$

Proof of Theorem 1.1. By Theorems 2.1, 3.1 and 3.2,

$$C_{\text{LO}} \leq \frac{3}{2} \left(2 \cdot 1.8150 \cdot \left(\frac{3}{5} \right)^2 \right)^{1/3}.$$

The right side is < 1.64 , proving the theorem. \square

4 What would be needed to beat 1.58

The same variational formula can produce 1.58 by optimizing over nontrivial radial measures μ, ν . A proof at the level of rigor of the Jellium/UEG papers would require the following finite certificate:

explicit rational radial measures μ, ν , an explicit rational piecewise upper function f ,

with

$$\Psi_{\mu\nu}(a, b) \leq f(a) + f(b) \quad \text{for all } a, b \geq 0,$$

and

$$\frac{3}{2} \left(2D(\mu, \mu)^2 \int_{\mathbb{R}^3} \frac{f(|z|)}{|z|^7} dz \right)^{1/3} < 1.58.$$

Without such a certificate, a smaller number is only a numerical experiment, not a proof of a new Lieb–Oxford constant.

Acknowledgements. LaTeX typesetting and bibliographic copy-editing were carried out with assistance from Claude Opus 4.7 (Anthropic). All mathematical content and proofs are the author’s own.

References

- [1] E. H. Lieb and S. Oxford. Improved lower bound on the indirect Coulomb energy. *Int. J. Quantum Chem.* **19** (1981), 427–439.
- [2] M. Lewin, E. H. Lieb, and R. Seiringer. Improved Lieb–Oxford bound and its application to density functional theory. *Pure Appl. Anal.* **2**, no. 1 (2020), 35–73.